

THE GENERATING FUNCTION OF TWO-STACK SORTABLE PERMUTATIONS BY DESCENTS IS REAL-ROOTED

PETTER BRÄNDÉN

ABSTRACT. Bóna has conjectured that, for fixed $n \geq 0$ and $t \geq 1$, the generating function of t -stack sortable permutations of length n by descents is real-rooted. The conjecture is known to be true for $t = 1$ and $t = n - 1$. Here we prove it for $t = 2$.

1. INTRODUCTION

Let $W_t(n, k)$ be the number of t -stack sortable permutations in the symmetric group, $\mathcal{S}_n$, with k descents. Recently, Bóna [1, 2] showed that for fixed n and t the numbers $W_t(n, k)$ form a unimodal sequence. Let $W_{n,t}(x) = \sum_{k=0}^{n-1} W_t(n, k)x^k$. A sufficient condition on a nonnegative sequence to be unimodal is that the corresponding polynomial is real-rooted. When $t = n - 1$ and $t = 1$ we get the Eulerian and the Narayana polynomials respectively. These are known to be real-rooted and Bóna conjectures that the same is true for general t . In what follows we will prove the conjecture for $t = 2$.

Let W be the set of finite words on $\mathbb{N}$ without repetitions. If w is any nonempty word we may write it as the concatenation $w = LnR$ where n is the greatest letter of w and L and R are the subwords to the left and right of n respectively. The stack-sorting operation $s : W \rightarrow W$ may be defined recursively by

$$s(w) = \begin{cases} w, & \text{if } w \text{ is the empty word,} \\ s(L)s(R)n, & \text{if } w = LnR \text{ is nonempty.} \end{cases}$$

The *stack sortable* permutations in $\mathcal{S}_n$ are the permutations which are mapped by s to the identity permutation. Similarly, a permutation is called *t -stack sortable* if $s^t(\pi)$ is the identity permutation.

2. STACK SORTABLE PERMUTATIONS AND JACOBI POLYNOMIALS

The number of stack sortable permutations of length n with k descents are known [10] to be the famous Narayana numbers [11, 12]

$$W_1(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k+1}.$$

The Narayana polynomials $W_{n,1}(x)$ are known to have real zeros. A simple proof of this fact is obtained by expressing $W_{n,1}(x)$ in terms of Jacobi

polynomials. Recall the definition of the *hypergeometric function* ${}_2F_1$:

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n z^n}{(c)_n n!},$$

where $(\alpha)_0 = 1$ and $(\alpha)_n = \alpha(\alpha+1) \cdots (\alpha+n-1)$ when $n \geq 1$. The Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$ can be expressed in the following two ways [9, Page 254]:

$$P_n^{(\alpha, \beta)}(x) = \frac{(1+\alpha)_n}{n!} {}_2F_1\left(-n, 1+\alpha+\beta+n; 1+\alpha; \frac{1-x}{2}\right), \quad (1)$$

$$P_n^{(\alpha, \beta)}(x) = \frac{(1+\alpha)_n}{n!} \left(\frac{x+1}{2}\right)^n {}_2F_1\left(-n, -\beta-n; 1+\alpha; \frac{x-1}{x+1}\right). \quad (2)$$

Rewriting $W_1(n+1, k)$ we end up with

$$W_{n+1,1}(x) = {}_2F_1(-n, -n-1; 2; x),$$

which by (2) gives

$$W_{n+1,1}(x) = \frac{1}{n+1} (1-x)^n P_n^{(1,1)}\left(\frac{1+x}{1-x}\right).$$

Since the Jacobi polynomials are orthogonal when $\alpha, \beta > -1$ we know that $W_{n,1}(x)$ is real- and simple-rooted and that the zeros of $W_{n,1}(x)$ strictly interlace the zeros of $W_{n+1,1}(x)$, that is, $\{W_{n+1,1}(x)\}_{n=0}^{\infty}$ form a *Sturm sequence*.

The numbers $W_2(n, k)$ are surprisingly hard to determine despite of their compact and simple form. It was recently shown that

$$W_2(n, k) = \frac{(n+k)!(2n-k-1)!}{(k+1)!(n-k)!(2k+1)!(2n-2k-1)!}.$$

See [3, 6, 7, 8] for proofs and more information on 2-stack sortable permutations.

A sequence of real numbers $\Gamma = \{\gamma_k\}_{k=0}^n$ is called an *n-sequence (of the first kind)* if for any real-rooted polynomial $f = a_0 + a_1x + \cdots + a_nx^n$ of degree at most n the polynomial $\Gamma(f) := a_0\gamma_0 + a_1\gamma_1x + \cdots + a_n\gamma_nx^n$ is real-rooted. There is a simple algebraic characterisation of *n-sequences* [5]:

Theorem 1. *Let $\Gamma = \{\gamma_k\}_{k=0}^n$ be a sequence of real numbers. Then Γ is an *n-sequence of the first kind* if and only if $\Gamma[(x+1)^n]$ is real-rooted with all its zeros of the same sign.*

We need the following lemma:

Lemma 2. *Let n be a positive integer and r a non-negative real number. Then $\Gamma = \left\{\binom{-n-r}{k}\right\}_{k=0}^n$ is an *n-sequence*.*

Proof. Let $r > 0$. Then

$$\begin{aligned} \Gamma[(x+1)^n] &= \sum_{k=0}^n \binom{-n-r}{k} \binom{n}{k} x^k \\ &= {}_2F_1(-n, n+r; 1; x) \\ &= P_n^{(0, r-1)}(1-2x), \end{aligned}$$

where the last equality follows from (1). Since the Jacobi polynomials are known to have all their zeros in $[-1, 1]$ when $\alpha, \beta > -1$ we have that $\Gamma[(x+1)^n]$ has all its zeros in $[0, 1]$. The case $r = 0$ follows by continuity when we let r tend to zero from above. $\square$

From the case $r = 0$ in Lemma 2 and the identity

$$\sum_{k=0}^n \binom{2n-k-1}{n-1} \binom{n}{k} x^k = (-1)^n \sum_{k=0}^n \binom{-n}{k} \binom{n}{k} (-x)^{n-k},$$

it follows that $\binom{2n-k-1}{n-1}$ is an n -sequence.

Theorem 3. *For all $n \geq 0$ the polynomial $W_{2,n}(x)$, which records 2-stack sortable permutations by descents, is real-rooted.*

Proof. We may write $W_2(n, k)$ as

$$W_2(n, k) = \frac{\binom{2n-k-1}{n-1} \binom{n+k}{n-1} \binom{2n}{2k+1}}{n^2 \binom{2n}{n}}.$$

A well known result on real-rooted polynomials reads as follows: If $\sum_i a_i x^i$ is a polynomial having only real non-positive zeros then so is the polynomial $\sum_i a_k i x^i$, where k is any positive integer. For a proof see [4, Theorem 3.5.4]. Applying this result to the polynomial $x(1+x)^{2n}$ we see that $\sum_k \binom{2n}{2k+1} x^k$ is real-rooted. Now,

$$\sum_{k=0}^{n-1} \binom{n+k}{n-1} \binom{2n}{2k+1} x^k = \sum_{k=0}^{n-1} \binom{2n-k-1}{n-1} \binom{2n}{2k+1} x^{n-1-k},$$

which by the discussion after Lemma 2 is real-rooted. Another application of Lemma 2 gives that $W_{n,2}(x)$ is real-rooted. $\square$

REFERENCES

- [1] Miklós Bóna. Corrigendum: “Symmetry and unimodality in t -stack sortable permutations” [J. Combin. Theory Ser. A **98** (2002), no. 1, 201–209; 1897934]. *J. Combin. Theory Ser. A*, 99(1):191–194, 2002.
- [2] Miklós Bóna. Symmetry and unimodality in t -stack sortable permutations. *J. Combin. Theory Ser. A*, 98(1):201–209, 2002.
- [3] M. Bousquet-Mélou. Multi-statistic enumeration of two-stack sortable permutations. *Electron. J. Combin.*, 5(1):Research Paper 21, 12 pp. (electronic), 1998.
- [4] F. Brenti. Unimodal, log-concave and Pólya frequency sequences in combinatorics. *Mem. Amer. Math. Soc.*, 81(413):viii+106, 1989.
- [5] T. Craven and G. Csordas. Multiplier sequences for fields. *Illinois J. Math.*, 21(4):801–817, 1977.
- [6] S. Dulucq, S. Gire, and O. Guibert. A combinatorial proof of J. West’s conjecture. *Discrete Math.*, 187(1-3):71–96, 1998.
- [7] I. P. Goulden and J. West. Raney paths and a combinatorial relationship between rooted nonseparable planar maps and two-stack-sortable permutations. *J. Combin. Theory Ser. A*, 75(2):220–242, 1996.
- [8] B. Jacquard and G. Schaeffer. A bijective census of nonseparable planar maps. *J. Combin. Theory Ser. A*, 83(1):1–20, 1998.
- [9] E. D. Rainville. *Special functions*. The Macmillan Co., New York, 1960.
- [10] R. Simion. Combinatorial statistics on noncrossing partitions. *J. Combin. Theory Ser. A*, 66(2):270–301, 1994.

- [11] Richard P. Stanley. *Enumerative combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999.
- [12] R. A. Sulanke. The Narayana distribution. *J. Statist. Plann. Inference*, 101(1-2):311–326, 2002. Special issue on lattice path combinatorics and applications (Vienna, 1998).

MATEMATIK, CHALMERS TEKNISKA HÖGSKOLA OCH GÖTEBORGS UNIVERSITET,
S-412 96 GÖTEBORG, SWEDEN
E-mail address: `branden@math.chalmers.se`